

Intratumoral *Fusobacterium* as prognostic factor in early stage colorectal cancer: results of the FUSOMAP study.

P. Nuciforo¹, F. Moratalla Navarro², I. Baraiibar Argota³, L. Alonso⁴, G. Serna⁵, J. Vidal Barrull⁶, L. Aguilera Castro⁷, N. Mulet Margalef⁸, A. Fernandez Montes⁹, M. Cuatrecasas⁶, I. Gardeazabal Gonzalez¹⁰, E. Gonzalez¹⁰, P. Garcia Alfonso¹¹, S. Rosello Keranen¹², R. Comas¹³, N. Bayo Puxan¹⁴, R. Fasanl¹⁵, M. Obon¹⁶, M.E. Elez Fernandez¹⁶, V. Moreno Aguado¹⁶

(1) Molecular Oncology Dept., Vall d'Hebron Institute of Oncology - Cellex Center, Barcelona, Spain
(2) Programa d'Atenció de Diagnòstic Oncològic (PADO) ICO - Institut Català d'Oncologia Hospital Duran i Reynald, Hospital de Llobregat, Spain
(3) Oncology Department, Vall d'Hebron University Hospital, Barcelona, Spain
(4) Medical Oncology Department, Hospital del Mar Research Institute, Universitat Pompeu Fabra, IDIBRON, Barcelona, Spain
(5) Gastroenterology, Vall d'Hebron University Hospital, Barcelona, Spain
(6) Medical Oncology, ICO - Institut Català d'Oncologia Badalona Hospital Universitari Germans Trias i Pujol, Badalona, Spain

(7) Dept. Medical Oncology, Complejo Hospitalario Universitario de Ourense, Ourense, Spain
(8) Pathology Hospital Clinic y Provincial de Barcelona, Barcelona, Spain
(9) Medical Oncology Department, HJMV - Hospital Universitario Marques de Valdecarlos, Santander, Spain
(10) Medical Oncology, Hospital Universitario Virgen de las Nieves, Granada, Spain
(11) Medical Oncology, Hospital Universitario Gregorio Marañón, Universidad Complutense, Madrid, Spain
(12) Medical Oncology, Hospital Clínico Universitario de Valencia, Valencia, Spain

BACKGROUND

The prevalence, geographical distribution, and prognostic significance of intratumoral *Fusobacterium* (iFUSO) infection in early stage colorectal cancer (CRC) is largely unknown. Here, we present the first results of a large national retrospective-prospective multicenter study (FUSOMAP) which aims to enhance our understanding on the pathophysiology of the CRC microbiome for the development of microbiota-based diagnostic and prognostic methods.


METHODS

Patients from different geographic and socioeconomic populations diagnosed at 9 hospitals across Spain as part of regional CRC screening programs were included in the present retrospective analysis. Formalin-fixed paraffin-embedded primary tumor tissue samples were retrieved for iFUSO quantification and visualization using an in-house developed RNA in situ hybridization assay. Positivity was defined as tumors with density

values ≥ 4 bacteria/mm². Cox proportional hazards model of disease-free survival (DFS) was used to compute hazard ratio (HR), controlling for potential confounders including gender, age, site, stage, grade, vascular invasion and adjuvant treatment.


RESULTS

1. Fusomap study patient cohort



2. Intratumoral *Fusobacterium* (iFUSO) prevalence

iFUSO was found in 21% of 700 evaluable CRC patients. The bacterium predominantly accumulates in the luminal surface in correspondence of the tumor (Figure 2A). The absolute amount of iFUSO varied significantly according to geography of primary tumor origin even after adjusting for confounding covariates (Figure 2B).




When classifying tumors according to iFUSO status, positivity rates ranged from 15.3% to 36.7% (from 5.9 to 22.3% within Catalunya, Figure 3).



3. Association between Intratumoral *Fusobacterium* (iFUSO) and clinicopathological characteristics


iFUSO positivity increased with tumor stage (Stage I-III: 1-26%; III-27%), tumor size, presence of vascular invasion, and decreased from right colon (27%) to rectum (16%) location (Table 1, Figure 4).



4. Association between Intratumoral *Fusobacterium* (iFUSO) and disease-free survival in stage I-III CRC


With a median follow-up of 3.76 years, patients with iFUSO-positive tumors had a significantly worse DFS compared to iFUSO-negative tumors (HR: 0.3, 95% CI: 0.13-0.68, p=0.007) (Figure 5).

In multivariable analysis, iFUSO remained significantly independently associated with DFS (p=0.04) when adjusted for age, sex, stage, grade, site, adjuvant chemotherapy and venous invasion. 3-years and 5-years DFS HRs (95% CI) for iFUSO-positive vs negative tumors were 1.68 (1.00-2.50) and 1.68 (1.02-2.4), respectively (Figure 6).



5. Association between Intratumoral *Fusobacterium* (iFUSO) and disease-free survival in patients with stages II-III CRC treated with adjuvant 5-FU-based chemotherapy.

For chemo-treated patients (39.4 % of the cohort), a statistically significant interaction was observed for the predictive value of iFUSO in term of DFS. iFUSO positivity significantly predicted worse outcome after standard-of-care 5-FU chemotherapy (HR: 1.85, CI: 1.11-3.07, p-value=0.018, Figure 7A). Conversely, no significant difference in DFS were observed according to iFUSO status in untreated patients (Figure 7B).



CONCLUSIONS

- iFUSO was found in 21% of stages I-III CRC patients.
- The absolute amount of iFUSO varied significantly according to geography of primary tumor origin even after adjusting for confounding covariates.
- iFUSO positivity increased with tumor stage, tumor size, presence of vascular invasion and decreased from right colon to rectum location.
- iFUSO was an independent negative prognostic factor in the overall population (stages I-III and in stages II-III patients treated with 5-FU adjuvant chemotherapy).