

BACKGROUND

The GEICO-88R study assessed the real-world use of niraparib as maintenance treatment in patients with PSROC (Cueva *et al*, *EJC* 2023), within the Spanish expanded access programme [1]. A preplanned subanalysis of long-term responders (niraparib exposure ≥ 12 months) was also reported (Cueva *et al*, *ESMO* 2023, 795-P) [2]. And, more recently, an additional subanalysis with extended follow-up was communicated in this population (Cueva *et al*, *ESMO GYN* 2024, 55P) [3].

OBJECTIVE

The goal of this work was to perform a post-hoc subanalysis of sustained long-term responders (SLTR); patients with niraparib exposure ≥ 24 months.


METHODS

In this retrospective study across 57 Spanish sites, patients received niraparib at a fixed 300 mg/day starting dose (FSD) or an individualized starting dose (ISD), 200 or 300 mg/day, according to weight and platelet count. In the current analysis a specific assessment of SLTR was performed, focusing on patient characteristics, niraparib dosing, effectiveness, tolerability, and post-niraparib systemic treatments. The results are presented in the poster comparing both the SLTR and non-SLTR (NSLTR) populations.

RESULTS

- Of 316 patients (general study), 61 (19.3%) SLTR were analyzed. In addition, NSLTR patients were also assessed.
- 82% were BRCAwt, with median of 2 prior treatment lines and relevant comorbidities (Table 1).
- Median niraparib initial dose was 200 mg/day. FSD/ISD was used in 16 (26.2%) and 45 (73.8%) patients, respectively. 37.7% of patients discontinued niraparib due to progression, 3.3% toxicity and 4.9% because of physician/patient decision (Table 2).
- The most frequent all-grade adverse events were thrombocytopenia, anemia and asthenia (fatigue). One acute myeloid leukemia (AML) was observed (Figure 1).
- 69.2% overall response rate (ORR) was achieved in the 26 patients with assessable pre-niraparib measurable disease; vs 20.6% in 116 NSLTR. With a median follow-up of 57 months, mPFS was 47.3 months (95% CI 28.8-NA). mPFS2 and mOS were not reached (Figures 2, 3, 4).
- 27 patients (44.3%) had ≥ 1 post-niraparib systemic treatment, in 23 patients (85.1%) was platinum-based (Table 3). ORR with the first subsequent line was 33.3%. mPFS of first post-niraparib line was 7.8 months (95% CI 3.6-17.5) vs 7.4 months (95% CI 6.2-8.1) in the NSLTR group.
- Upon analysis, 31 (50.8%) patients remained on niraparib therapy, and 42 (68.9%) patients were still alive, 15 (24.6%) of them without apparent illness. 56 (91.8%) and 41 (67.2%) patients were alive at 3 and 5 years, respectively.

Figure 1. Niraparib toxicities



REFERENCES

1. Real-world safety and effectiveness of maintenance niraparib for platinum-sensitive recurrent ovarian cancer: A GEICO retrospective observational study within the Spanish expanded-access programme. *European Journal of Cancer*, January 2023.
2. 795P – Tolerability and effectiveness of niraparib in long-term responders with platinum-sensitive recurrent ovarian cancer (GEICO-88R study). Poster presentation. *ESMO* 2023.
3. 55P – Long-term responders (LTR) with niraparib maintenance in platinum-sensitive recurrent high-grade serous ovarian cancer (PSROC) focusing on subsequent therapies and post-progression outcomes (GEICO-88R study). Poster presentation. *ESMO GYN* 2024.

Table 1. Patient characteristics

		SLTR (N=61)	NSLTR (N=255)
Median age		61 (45-80)	64 (31-88)
Most common initial FIGO	IIIC	36 (59.0%)*	122 (47.8%)*
	IVA	4 (6.6%)	31 (12.2%)
	IVB	0 (0.0%)	43 (16.9%)
BRCA status	BRCAmut	4 (6.6%)	14 (5.5%)
	BRCAwt	50 (82.0%)	204 (80.0%)
	Unknown	7 (11.5%)	37 (14.5%)
Median previous chemotherapy lines		2 (1-8)	3 (1-11)
	1	1 (1.6%)	5 (2.0%)
	2-4	57 (93.4%)	207 (81.1%)
	>4	3 (4.9%)	43 (16.9%)
Previous bevacizumab and PARPI	Bevacizumab	24 (39.3%)	118 (46.2%)*
	PARPI	2 (3.3%)	21 (8.2%)*
Surgery after relapse	R0	15 (71.4%)	33 (63.5%)*
	R>0	4 (19.0%)	13 (25.0%)*
	Unknown	2 (9.5%)	6 (11.5%)
ECOG	0	36 (62.3%)*	121 (47.5%)*
	1	23 (37.7%)*	127 (49.8%)*
	Unknown	0 (0.0%)	7 (2.7%)
Pre-niraparib measurable and assessable disease**	Yes	26 (42.6%)*	116 (45.5%)*
	No	35 (57.7%)*	139 (54.5%)*
Most common relevant comorbidities		29 (47.5%)*	100 (39.2%)*
Arterial hypertension		13 (44.8%)*	62 (62.0%)*
Diabetes mellitus		3 (10.3%)*	21 (21.0%)*
Obesity		4 (13.8%)*	8 (8.0%)*

* Results with bold letters indicate significant (>10% difference) relevance.

** From patients with measurable disease before niraparib, ORR was calculated in those having assessable RECIST 1.1 response.

Table 3. First and second lines after niraparib

PBT: Platinum-based treatment. PRS: Platinum-resistant scheme (other than paclitaxel).

	SLTR (N = 61) - N (%)	NSLTR (N = 255) - N (%)
1st line after niraparib		
PBT + bevacizumab	5 (18.5)	24 (11.9)
PBT	11 (40.7)	85 (33.3)
Paclitaxel + bevacizumab	--	25 (9.8)
Paclitaxel	--	10 (3.9)
PRS + bevacizumab	--	8 (3.1)
PRS	--	11 (4.3)
PARPI	--	4 (1.5)
Other	11 (40.7)	34 (13.3)
2nd line after niraparib		
PBT + bevacizumab	1 (5.2)	--
PBT	6 (31.5)	27 (18.4)
Paclitaxel + bevacizumab	2 (10.5)	17 (11.6)
Paclitaxel	2 (10.5)	25 (17.1)
PRS + bevacizumab	1 (5.2)	4 (2.7)
PRS	2 (10.5)	17 (11.6)
PARPI	1 (5.2)	5 (3.4)
Other	4 (20.8)	51 (34.9)

CONCLUSIONS

This subanalysis of SLTR to niraparib maintenance in real life shows a significant proportion of patients (nearly 20% of the total) with remarkable benefit (≥ 24 months), some of them without detectable tumour. Two thirds of patients are alive at 5 years and mOS is not reached despite being a high-risk population. Most of them received a platinum-based treatment as next systemic line after niraparib progression. Although it is impossible to identify significant predictors from a retrospective descriptive study, in addition to better tolerability and niraparib response, there are relevant differences between the SLTR and NSLTR populations (highlighted in bold) that could explain the divergent outcomes.

Table 2. Treatment disposition

		SLTR (N=61)	NSLTR (N=255)
Initial dose (mg)	200	38 (62.3%)*	140 (54.9%)*
	300	23 (37.7%)*	115 (45.1%)*
Individualized	Yes (ISD)	45 (73.8%)*	194 (76.1%)*
	No (FSD)	16 (26.2%)*	61 (23.9%)*
Median dose (mg)		200	200
Treatment duration (months)	Median	57.2 (24.4-77.3)	5.6 (0.2-22.5)
	0-12	0 (0.0%)*	211 (82.7%)*
	12-24	0 (0.0%)*	44 (17.3%)*
	>24	61 (100.0%)*	0 (0.0%)*
Dose interruptions		46 (75.4%)*	154 (60.3%)*
Dose reductions		46 (75.4%)*	142 (55.6%)*
Still on treatment upon analysis		31 (50.8%)*	0 (0.0%)*
End of treatment (other than progression)	Toxicity	2 (3.3%)*	15 (5.8%)*
	Physician's decision	1 (1.6%)*	7 (2.7%)*
	Patient's decision	2 (3.3%)*	4 (1.6%)*
	Other	2 (3.3%)*	5 (1.9%)*

Figure 2. Progression-free survival (47.3 mo SLTR vs 6.2 mo non-SLTR)




Figure 3. Progression-free survival 2 (NA SLTR vs 16.5 mo non-SLTR)





Figure 4. Overall survival (NA SLTR vs 26.9 mo non-SLTR)



ACKNOWLEDGEMENTS

GEICO would like to thank all the patients who contributed to this study along with their families. Special thanks to all participating investigators. Funding for this study was provided by GSK (NCT04546373). GSK was given the opportunity to perform a courtesy review of the preliminary version of this poster for accuracy only, but the authors are solely responsible for final content and interpretation.