

PREDICTION OF CLINICALLY SIGNIFICANT BLEEDING IN ANTICOAGULATED PATIENTS FOR CANCER-ASSOCIATED VENOUS THROMBOEMBOLISM: VALIDATION OF THE B-CAT SCORE IN A COHORT FROM THE TESEO STUDY

J. López Robles¹, M. Sánchez Cánovas¹, C. Iglesias², L. Ortega Morán³, L. Sánchez Togneri⁴, J. Rubio⁵, I. Fernández Pérez⁶, M. García De Herreros⁷, I. García Escobar⁸, R. Porta⁹, E. Brozos¹⁰, M. Carmona Campos¹¹, E. Martínez De Castro¹², H. Olivares¹³, T. Quintanar¹⁴, S. García Adrián³, M. Covela Rúa¹⁵, A. Carmona Bayonas¹, P. Jiménez Fonseca², AJ. Muñoz Martín³, On Behalf Of Teseo Registry Investigators.

1. Medical Oncology Department, Hospital Universitario Morales Meseguer, University of Murcia, IMB, Murcia, Spain; 2. Medical Oncology Department, Hospital Universitario Central de Asturias, ISPA, University of Oviedo, Oviedo, Spain; 3. Medical Oncology Department, Hospital General Universitario Gregorio Marañón, Madrid, Spain; 4. Medical Oncology Department, Hospital Universitario de Basurto, Bilbao, Spain; 5. Medical Oncology Department, Hospital Universitario Fundación Jiménez Díaz, Madrid, Spain; 6. Medical Oncology Department, Hospital Álvaro Cunqueiro, Complejo Hospitalario Universitario de Vigo, Vigo, Spain; 7. Medical Oncology Department, Hospital Clínic de Barcelona, Barcelona, Spain; 8. Medical Oncology Department, Hospital Universitario de Ciudad Real, Ciudad Real, Spain; 9. Medical Oncology Department, Hospital Universitari Dr. Josep Trueta, Institut Catalán de Oncologia, Girona, Spain; 10. Medical Oncology Department, Hospital Universitario de A Coruña, A Coruña, Spain; 11. Medical Oncology Department, Complejo Hospitalario Universitario de Santiago, Santiago de Compostela, Spain; 12. Medical Oncology Department, Hospital Universitario Marqués de Valdeoln, Instituto de Investigación IDIVAL, Santander, Spain; 13. Medical Oncology Department, Hospital Universitario 12 de Octubre, Madrid, Spain; 14. Medical Oncology Department, Hospital General Universitario de Elche, Elche, Spain; 15. Medical Oncology Department, Hospital Universitario Lucas Augusti, Lugo, Spain.

BACKGROUND

There is a lack of a validated scoring system for predicting clinically significant bleeding in patients anticoagulated for cancer-associated venous thromboembolism (Ca-VTE).

OBJECTIVES

The aim was to validate the B-CAT score, a new tool designed to classify the risk of bleeding in oncology patients receiving anticoagulation.

METHODS


Data came from the TESEO study, a national, multicenter and prospective registry that documents patients with Ca-VTE. Patients anticoagulated were included and observed for 180 days for major or clinically relevant bleeding. Variables of the B-CAT score were selected, except for minor trauma, and minor surgery and clinically relevant bleeding not requiring hospitalization after Ca-VTE, as these were not available. Patients were classified according to bleeding risk into three categories, and a multivariate logistic regression model was developed to estimate the risk of bleeding.

RESULTS

BASELINE CHARACTERISTICS (N = 2,301)

		TESEO (n = 2,301)	Cohen et al. (n = 15,749)
Sex	Male	1,163 (50.1%)	7,625 (48.4%)
Age	Median (p25, p75)	65 (58-73)	68 (60-75)
ECOG PS	0 - 1	1,679 (72.9%)	-
	2 - 4	622 (27.1%)	-
Cancer type	Breast	244 (10.6%)	2,906 (18.5%)
	Cervix	29 (1.3%)	144 (0.9%)
	Uterus	51 (2.2%)	379 (2.4%)
	Ovarian	99 (4.3%)	562 (3.6%)
	Prostate	51 (2.2%)	1,639 (10.4%)
	Central nervous system	62 (2.7%)	418 (2.7%)
	Upper gastrointestinal tract	154 (6.7%)	994 (6.3%)
	Colorectal and other low gastrointestinal tract	436 (18.9%)	2,605 (16.5%)
	Hepatopancreatobiliary	292 (12.7%)	677 (4.3%)
	Lung	523 (22.8%)	1,538 (9.8%)
	Kidney	36 (1.6%)	354 (2.2%)
	Bladder	82 (3.6%)	647 (4.1%)
	Hematologic	11 (0.5%)	1,418 (9%)
	Malignant melanoma	22 (1%)	260 (1.7%)
	Head and neck	51 (2.2%)	-
Cancer therapy	Other	148 (6.4%)	915 (5.8%)
	Unknown primary site	10 (0.4%)	293 (1.9%)
	Metastases	1,628 (70.8%)	7,052 (44.8%)
	None	76 (3.3%)	3,831 (24.3%)
	Chemotherapy	1,044 (45.4%)	7,905 (50.2%)
	Combination chemotherapy	301 (13.1%)	-
	Targeted therapy	79 (3.4%)	-
	Immunotherapy	97 (4.2%)	-
	Hormonal	55 (2.4%)	2,064 (13.1%)
	Other	55 (2.4%)	-
Venous thromboembolism	Radiation	599 (26%)	1,949 (12.4%)
	Deep vein thrombosis of the legs	585 (25.4%)	6,835 (43.4%)
	Pulmonary embolism	1,296 (56.3%)	8,914 (56.6%)
	Other	612 (26.6%)	-
Anticoagulant treatment	Low-molecular-weight heparin	2,238 (97.2%)	10,820 (68.7%)
	Direct Oral Anticoagulant	50 (2.2%)	2,039 (12.9%)
	Vitamin K antagonist	13 (0.6%)	2,890 (18.4%)

CLINICALLY SIGNIFICANT BLEEDING



PREDICTORS FOR SIGNIFICANT BLEEDING


Cancer type	436 (18.9%)
Metastases	1,628 (70.8%)
Minor surgery after Ca-VTE	Not available
Minor trauma	Not available
Major bleeding before Ca-VTE	15 (0.7%)
CRNMB < 731 days before Ca-VTE	31 (1.3%)
CRNMB not requiring hospitalization after Ca-VTE	Not available
Stroke	75 (3.3%)
Gastroduodenal disease	189 (8.2%)
Anaemia	1,216 (52.8%)
Coagulation disorder	20 (0.9%)

BLEEDING RISK ACCORDING TO B-CAT

Score category	Bleeding events, n (%)	Person-years	Incidence rate, events/100 p-y (95% CI)	Cohen et al.
Low-risk group (n=1,101)	51 (4.6%)	449	11.4 (8.3–14.5)	5.12 (4.4–6)
Medium-risk group (n=1,161)	95 (8.2%)	389	24.4 (19.5–29.3)	18.96 (6.9–21.2)
High-risk group (n=39)	11 (28.2%)	11	100 (41–159)	58.54 (44–71.6)


p < 0.001

PREDICTIVE MODEL: DISCRIMINATION



	C-statistic TESEO	Cohen et al.
Complete cohort	0.627 (0.578 – 0.675)	0.7 (0.65 – 0.75)
Major bleeding	0.609 (0.531 – 0.687)	0.76 (0.68 – 0.84)
CRNMB-H	0.632 (0.571 – 0.692)	0.67 (0.61 – 0.73)

PREDICTIVE MODEL: CALIBRATION



CONCLUSIONS

We have validated the bleeding risk score B-CAT in patients with Ca-VTE receiving anticoagulation. This model can contribute to standardizing decision-making in a context where quality evidence is limited.